



A practical guide for staff

Managing infectious cases of infectious in all education and childcare settings

The online guide offers information and resources for the following:

Introduction to infection control in all education and childcare settings

Prevention and control

Outbreak management

Immunisation

Staff health, pet and animal contact

Cleaning the environment

Managing specific infections

Exclusion table

Diarrhoea and vomiting outbreak action list



Advice and guidance

To find out more, visit www.gov.uk/government/publications/ health-protection-in-schools-and-other-childcare-facilities or bit.ly/2xiJpPZ.





Should I keep my

child off school?



Until...

Chickenpox	at least 5 days from the onset of the rash and until all blisters have crusted over
Diarrhoea and Vomiting	48 hours after their last episode
Cold and Flu-like illness (including COVID-19)	they no longer have a high temperature and feel well enough to attend. Follow the national guidance if they've tested positive for COVID-19.
Impetigo	their sores have crusted and healed, or 48 hours after they started antibiotics
Measles	4 days after the rash first appeared
Mumps	5 days after the swelling started
Scabies	they've had their first treatment
Scarlet Fever	24 hours after they started taking antibiotics
Whooping Cough	48 hours after they started taking antibiotics

No

but make sure you let their school or nursery know about...

Hand, foot and mouth	Glandular fever
Head lice	Tonsillitis
Threadworms	Slapped cheek





Advice and guidance

To find out more, search for health protection in schools or scan the QR code or visit https://qrco.de/minfec.